arXiv:1504.05370v3 [hep-th] 19 Aug 2015

Notes on stress energy tensor of neutral blackfolds

J. Sadeghi[a,1], Z.Amoozad,[a,*]

[a] *Department of physics, University of Mazandaran,*
Babolsar, Iran

Abstract

In this paper we review some properties of higher dimensional black holes. We take advantages of fluid/gravity duality and obtain the fluid properties of higher dimensional black holes on the boundary. So we consider two neutral and charged blackfold cases and extract the Brown-York stress energy tensor of fluid. But Brown-York method does not work for all intrinsic metric in the reference spacetime. Surprisingly, in AAds spacetimes the expectation value of the stress tensor in the CFT side solves the problem. In our case the neutral blackfold's spacetime is Ricci-flat. As we know, the compactification of some directions in any asymptotically AdS black brane corresponds to a Ricci-flat spacetime. So by calculating the AAdS form of that metric the dual renormalized holographic stress tensor has been extracted. This stress tensor is conserved and traceless, also it is same as perfect fluid.

Keywords: `Fluid/Gravity duality; Blackfold; Boundary stress tensor`

---

1. Introduction

Recently a nice relation between fluid and gravity has been emerged. This duality comes from the gauge/gravity duality which gives an equivalence between D dimensional gravitational theory and D-1 dimensional quantum field theory. It connects asymptotically AdS geometry to the states in the field the-

---

[*] Corresponding author, z.amoozad@stu.umz.ac.ir
[1] pouriya@ipm.ac.ir

ory side. The pure AdS bulk geometry corresponds to the vacuum state of the gauge theory. But deformed bulk geometry, in the way that AdS asymptotic properties preserves, could be corresponded to an exited state. The perturbed metric on the gravity side corresponds to the expectation value of stress energy tensor on the CFT side. In general this is the holographic principle that first has been proposed by Gerard 't Hooft in 1993, which is one of the most important subjects that is originated from the study of quantum gravity and string theory [1, 2, 3, 4, 5].

Near thermal equilibrium, quantum field theory describes the existence of a hydrodynamic regime that shows the long wavelength behavior of the microscopic degrees of freedom. So by acceptance of such holography, a hydrodynamic phenomenon can be reproduced and also the existence of fluid/gravity duality is proven [6, 7, 8].

One of the main difference of higher dimensional black holes with respect to the 4-dimensional ones is that the horizons of higher dimensional black holes can have at least two very different size lengths. By definition, if black holes regarded as a black brane (possibly boosted locally) which folded into multiple dimensions a blackfold is shaped. One reason that blackfolds can have at least two very different size horizon is that there is no bound on their angular momentum, although in 4-dimension the Kerr bound $J \leq GM^2$ does not admit higher momentum. So in blackfolds, there is two length scales which are given by,

$$l_M \sim (GM)^{1/D-3} \qquad , \qquad l_J \sim \frac{J}{M} \tag{1}$$

In case which $l_J \gg l_M$, this separation suggests an effective description of long wavelength dynamics [9]. Also it may causes to the Gregory-Laflamme instability of blackfolds. One of the simplest examples of the theory of blackfolds which defines an effective theory is a black ring (as a boosted black string) [10].

As the theory of classical brane dynamics [11] explains a long wavelength effective theory, black branes can also take the form of a dynamic fluids that lives on a dynamical worldvolume. But if higher dimensional black holes regarded instead of the 4-dimensional ones, then how we can extend the fluid/gravity correspondence? The holographic duality for Ricci-flat spacetime is one of the interesting dualities which recently people has been worked on.

Many different subjects have been investigated in these grounds. For example, AdS black brane solutions correspond to conformal and non conformal fluid at thermal equilibrium [12, 13] and also some relativistic expansion of fluid/gravity duality have been studied in [14, 15]. Extension of fluid/gravity duality to spherical horizon topologies, de sitter spacetime and charged fluid are calculated in [16, 17, 18, 19, 20, 21, 22].

Using the AdS/Ricci-flat correspondence and neutral blackfold as a Ricci-flat case, we extract AAdS form of a neutral blackfold in the Fefferman-Graham coordinate and the dual renormalized stress tensor which is a property of the boundary have been obtained. This paper is organized as follows. In the next section some properties and metric of the neutral and charged blackfolds have been reviewed. In section 3 the Brown-York stress energy tensor of blackfolds on a time-like hypersurface is extracted. Section 4 is dedicated to the calculation of stress tensor of the dual theory. In final section we conclude and note some points.

2. General properties of blackfolds

2.1. Neutral blackfolds

As we know 4-dimensional black holes only have short scale dynamics ($\sim r_0$). But for higher dimensional black holes we see some more physical information and have at least two very different size horizons $r_0 \ll R$. This properties predicts long distance worldvolume effective theory. So for higher dimensional ones new methods are needed to capture the long distance $\sim R \gg r_0$ dynamics [9]. In general, brane-like objects are sources of energy momentum tensor [23]. The requirement of consistent coupling of source to gravity overtakes,

$$\nabla_\mu T^{\mu\nu} = 0 \qquad \longrightarrow \qquad T^{\mu\nu} K_{\mu\nu}{}^{\rho} = 0 \tag{2}$$

where μ, ν are spacetime coordinates $(\mu, \nu = 0, ..., D-1)$ and ρ is the orthogonal one. ${K_{\mu\nu}}^{\rho}$ is the extrinsic curvature tensor of the blackfold embedding. On scales $r \ll R$, the blackfold is equivalent to a black p-brane up to a Lorentz transformation which can depend on position. The effective stress tensor of a static neutral flat black brane with orthonormal worldvolume coordinates (if the velocity field is u^a with $u^a u^b \eta_{ab} = -1$ *and* $\sigma^a = (\sigma^0, \sigma^i)$) is given by [9]:

$$T^{ab} = r_0^n (n u^a u^b - \eta^{ab}) \tag{3}$$

where σ^a is the worldvolume coordinates and $a, b = 0, ..., p$. As mentioned before, the blackfolds can have $\frac{l_M}{l_J} \to 0$, so that a flat black brane produces, and effective theory describes the collective dynamics of a black p-brane where it's geometry in $D = n + p + 3$ spacetime dimension is [11],

$$ds^2{}_{p-brane} = -(1 - \frac{{r_0}^n}{r^n})dt^2 + \sum_{i=1}^{p} d{z_i}^2 + (1 - \frac{{r_0}^n}{r^n})^{-1} dr^2 + r^2 d\Omega^2{}_{n+1} \tag{4}$$

where $\sigma^a = (t, z^i)$ is related to brane worldvolume. By boosting it along the worldvolume, one can easily find following metric,

$$ds^2{}_{p-brane} = (\eta_{ab} + \frac{{r_0}^n}{r^n} u_a u_b) d\sigma^a d\sigma^b + (1 - \frac{{r_0}^n}{r^n})^{-1} dr^2 + r^2 d\Omega^2{}_{n+1} \tag{5}$$

where r_0 is the thickness of the horizon. Here $(p+1)$ coordinates are on the worldvolume of the blackfold and $(D - p - 1)$ coordinates are transverse directions to the worldvolume. By taking long distance effective theory, the effective Einstein's equation can be written as,

$$R_{\mu\nu}{}^{(long)} - \frac{1}{2} R^{(long)} g_{\mu\nu}{}^{(long)} = 8\pi G T_{\mu\nu}{}^{(eff)}. \tag{6}$$

So the effective worldvolume stress tensor is,

$$T_{\mu\nu}{}^{(eff)} = -\frac{2}{\sqrt{-g_{(long)}}} \frac{\delta I_{eff}}{\delta g^{\mu\nu}{}_{(long)}} |_{W_{p+1}} . \tag{7}$$

As we know the quasilocal stress energy tensor is introduced by Brown and York [24]. It appropriately matches the coupling of the short wavelength and long wavelength degrees of freedom. If D_a be the covariant derivative with respect to the boundary metric γ_{ab}, the equations of conservation of the quasi local stress tensor will takes the following form,

$$D_a {T_{ab}}^{(quasilocal)} = 0 \tag{8}$$

This is the dynamics of effective fluid that lives on the worldvolume and spanned by the brane. For isotropic worldvolume theory, in case of lowest derivative order, the stress tensor will be the same as the isotropic perfect fluid, which is given by,

$$T^{ab} = (\varepsilon + p)u^a u^b + p\gamma^{ab} \tag{9}$$

where ε is the energy density and p is the pressure. If it doesn't be a perfect fluid then the stress tensor will have dissipative terms proportional to gradients of $\ln r_0, u^a, \gamma^{ab}$. For stationary spacetimes the effect of dissipative terms vanish. By definition $\overline{\nabla}_\mu = {h_\mu}^\nu \nabla_\nu$ where ∇_μ is the background covariant derivative and $h^{\mu\nu} = \partial_a X^\mu \partial_b X^\nu \gamma^{ab}$ and X^μ is the embedding function. Because of the consistent coupling of worldvolume to the long wavelength gravitational field $g_{\mu\nu}$ the stress tensor must obey,

$$\overline{\nabla}_\mu T^{\mu\rho} = 0 \tag{10}$$

The D equations in (10) separate into $D-p-1$ equations in directions orthogonal to W_{p+1} and $p+1$ equations parallel to W_{p+1}, which are given by,

$$\begin{aligned} T^{\mu\nu} {K_{\mu\nu}}^\rho = 0 \quad & \textit{extrinsic equation}\,, \\ D_a T^{ab} = 0 \quad & \textit{intrinsic equation}\,. \end{aligned} \tag{11}$$

2.2. Charged blackfolds

In supergravity and low energy limit of string theory, black branes have charges. So for higher dimensional cases, effective worldvolume theory for such charged branes is regarded. One of the best choices is black p-branes that carry charges of Ramond-Ramond field strength type $F_{(p+2)}$. The charged dilatonic black p-brane solution of action is [25],

$$I = \frac{1}{16\pi G} \int dx^D \sqrt{-g}(R - \frac{1}{2}(\partial\phi)^2 - \frac{1}{2(p+2)!} e^{a\phi} F^2_{(p+2)}) \tag{12}$$

where

$$a^2 = \frac{4}{N} - \frac{2(p+1)n}{D-2} \quad , \quad n = D - p - 3 \tag{13}$$

The flat black p-brane solution reads,

$$ds^2 = H^{\frac{-Nn}{D-2}}(-f dt^2 + \sum_{i=1}^{p} dz_i{}^2) + H^{\frac{N(p+1)}{D-2}}(f^{-1}dr^2 + r^2 d\Omega^2{}_{n+1}) \tag{14}$$

$$\begin{aligned} e^{2\phi} = H^{aN} \quad &, \quad A_{p+1} = \sqrt{N}\coth\alpha(H^{-1}-1)dt \wedge dz_1 \wedge dz_2 ... \wedge dz_p \, , \\ H = 1 + \frac{r_0{}^n \sinh^2\alpha}{r^n} \quad &, \quad f = 1 - \frac{r_0{}^n}{r^n}. \end{aligned} \tag{15}$$

where $\sinh\alpha$ related to the boost of the blackfold which is a Lorentz transformation of some of directions of p-brane. As $a^2 \leq 0$, the parameter N is not arbitrary and there is,

$$N \leq 2(\frac{1}{n} + \frac{1}{p+1}) \tag{16}$$

In string/M theory, N is an integer up to 3 (when $p \geq 1$)that corresponds to the number of different types of branes in an intersection. In this case the effective stress tensor can be written as,

$$T^{ab} = \tau s(u^a u^b - \frac{a}{b}\gamma^{ab}) - \Phi_p Q_p \gamma_{ab} \tag{17}$$

where τ is temperature and s is entropy and Φ_p is the potential that measured from the difference between the values of A_{p+1} at the horizon at $r \to \infty$ in (15). As one can see, the relation (17) surprisingly has a brane tension component $-\Phi_p Q_p$ and a thermal component τs. It must be noted that, blackfolds with p-brane charges do not have open boundaries; as the charge Q_p would not be conserved at boundaries. So it is possible to consider that blackfolds end on another brane that carries appropriate charge.

3. Brown-York stress energy tensor

3.1. Charged blackfolds

In the fluid/gravity duality, essential physical properties of field theory side are mentioned by boundary stress tensor. As we know the Brown-York stress energy tensor [26], which is induced by the bulk geometry, can be obtained by finding an appropriate hypersurface in the bulk. The bulk metric described by (14). Hypersurfaces can be obtained by constraining independent coordinates of spacetime. The time-like hypersurfaces play important role for causality relations between different coordinates. The hypersurface considered here is defined by:

$$r = r_c, \tag{18}$$

As we know, vectores perpendicular to a time-like hypersurface are space-like and connects time-like hypersurfaces. In that case, the Brown-York stress energy tensor [22] is defined by:

$$T_{ab} = 2(K\gamma_{ab} - K_{ab}), \tag{19}$$

where γ_{ab} is the induced metric on the time-like hypersurface and K_{ab} is the extrinsic curvature of the bulk. This can be determined by,

$$K_{\mu\nu} = -n_\mu a_\nu - n_{\nu;\mu} \ , \tag{20}$$

n_λ is the unit normal vector to the hypersurface,

$$n_\lambda = \frac{\phi_{,\lambda}}{|g^{\mu\nu}\phi_{,\mu}\phi_{,\nu}|^{\frac{1}{2}}} \ , \tag{21}$$

$$a_\nu = n^\theta n_{\nu;\theta} \ . \tag{22}$$

and ϕ in the (21) is the equation of hypersurface. In case $r = r_c$, ϕ just depends on r. The charged dilatonic blackfold (14) is,

$$ds^2{}_{p-brane} = Adt^2 + B{dz_i}^2 + Cdr^2 + F_\alpha d\psi^2{}_\alpha \tag{23}$$

and

$$\begin{aligned}
A &= -(1 + \frac{{r_0}^n \sinh^2\alpha}{r^n})^{\frac{-Nn}{D-2}}(1 - \frac{{r_0}^n}{r^n}),\\
B &= (1 + \frac{{r_0}^n \sinh^2\alpha}{r^n})^{\frac{-Nn}{D-2}},\\
C &= (1 + \frac{{r_0}^n \sinh^2\alpha}{r^n})^{\frac{N(p+1)}{D-2}}(1 - \frac{{r_0}^n}{r^n})^{-1},\\
F_\alpha &= F_\alpha(r, \psi_\alpha).
\end{aligned} \tag{24}$$

This metric is diagonal, so the inverse matrix can be found easily. For such a metric it is tried to determine the extrinsic curvature (20). In order to obtain the extrinsic curvature, one can arrange the n_λ, n^θ and a_ν as,

$$\begin{aligned} n_\lambda &= \sqrt{C}\delta_\lambda{}^r, \\ n^\theta &= \sqrt{C}g^{\theta r}, \\ a_\nu &= n^\theta(n_{\nu,\theta} - \Gamma^r{}_{\nu\theta}\sqrt{C}). \end{aligned} \tag{25}$$

So, by using the equation (25) into equation (20) one can obtain the corresponding extrinsic curvature as,

$$K_{\mu\nu} = -\delta_\mu{}^r\delta_\nu{}^r\frac{C_{,r}}{2\sqrt{C}} + C\sqrt{C}\delta_\mu{}^r g^{rr}\Gamma^r{}_{\nu r} - \frac{C_{,\mu}}{2\sqrt{C}}\delta_\nu{}^r + \sqrt{C}\Gamma^r{}_{\mu\nu} \tag{26}$$

By calculating the Christoffel symbols the nonvanishing components of extrinsic curvature for that geometry are given by,

$$K_{tt} = -\frac{A_{,r}}{2\sqrt{C}} \quad , \quad K_{ii} = -\frac{B_{,r}}{2\sqrt{C}} \quad , \quad K_{\psi_\alpha\psi_\alpha} = -\frac{F_{\alpha,r}}{2\sqrt{C}} \tag{27}$$

On the other side, the induced metric γ_{ab} on the time-like hypersurface is,

$$ds^2|_{r=r_c} = A_c dt^2 + B_c dz_i{}^2 + F_{\alpha c} d\psi^2{}_\alpha, \tag{28}$$

The trace of the extrinsic tensor can be written by following equation,

$$K = K_{ab}\gamma^{ab} = \frac{-1}{2\sqrt{C}}\{\frac{A_{,r}}{A_c} + p\frac{B_{,r}}{B_c} + \sum_{\alpha=1}^{n+1}\frac{F_{\alpha,r}}{F_{\alpha c}}\}, \tag{29}$$

where p is the number of spatial coordinates of worldvolume of blackfold. So the Brown-York stress energy tensor can easily be constructed. For the case of D/NS-brane in type II string theory [23] we have,

$$D = 10 \quad , \quad N = 1 \quad , \quad p = 0, ..., 6 \tag{30}$$

Following the definition,

$$\mathbf{A} = \frac{A_{,r}}{A_c} \quad , \quad \mathbf{B} = \frac{B_{,r}}{B_c} \quad , \quad \mathbf{F}_\alpha = \frac{F_{\alpha,r}}{F_{\alpha c}} \quad , \tag{31}$$

in case of $p = 5$ brane ($n = 2$), the nonvanishing component of Brown-York tensor on $r = r_c$ are,

$$T_{tt} = \frac{-1}{\sqrt{C}}\{[\mathbf{A} + 5\mathbf{B} + \mathbf{F}_1 + \mathbf{F}_2 + \mathbf{F}_3]A_c - A_{,r}\} \quad , \tag{32}$$

$$T_{ii} = \frac{-1}{\sqrt{C}}\{[\mathbf{A} + 5\mathbf{B} + \mathbf{F}_1 + \mathbf{F}_2 + \mathbf{F}_3]B_c - B_{,r}\} \quad , \tag{33}$$

$$T_{\alpha\alpha} = \frac{-1}{\sqrt{C}}\{[\mathbf{A} + 5\mathbf{B} + \mathbf{F}_1 + \mathbf{F}_2 + \mathbf{F}_3]F_{\alpha c} - F_{\alpha,r}\}. \tag{34}$$

One can easily check that there is a trace anomaly here.

3.2. Neutral blackfolds

The case of neutral blackfold has been reviewed in the section 2 and the appropriate metric found to be the equation (4). Here the method is the same as the last subsection, and we only present $p = 5$ brane which characterized by the metric,

$$ds^2 = Adt^2 + Bd{z_i}^2 + Cdr^2 + F_\alpha d\psi^2{}_\alpha \tag{35}$$

and

$$\begin{aligned} A &= -(1 - \frac{{r_0}^2}{r^2}), \\ B &= 1, \\ C &= (1 - \frac{{r_0}^2}{r^2})^{-1}, \\ F_\alpha &= F_\alpha(r, \psi_\alpha). \end{aligned} \tag{36}$$

The method is straightforward and the nonvanishing components of stress tensor for neutral blackfold at $r = r_c$ is,

$$T_{tt} = \frac{6}{r_c}(1 - \frac{{r_0}^2}{r_c^2})^{\frac{1}{2}}(1 + \frac{{r_0}^2}{r_c^2}), \tag{37}$$

$$T_{ii} = -(1 - \frac{{r_0}^2}{r_c^2})^{\frac{1}{2}}\{\frac{\frac{2{r_0}^2}{r_c^3}}{(1 - \frac{{r_0}^2}{{r_c}^2})} + \frac{6}{r_c}\}, \tag{38}$$

$$T_{\theta\theta} = -{r_c}^2(1-\frac{{r_0}^2}{r_c^2})^{\frac{1}{2}}\{\frac{\frac{2{r_0}^2}{r_c^3}}{(1-\frac{{r_0}^2}{{r_c}^2})}+\frac{4}{r_c}\}, \tag{39}$$

$$T_{\phi\phi} = \sin^2\theta \, T_{\theta\theta}, \tag{40}$$

$$T_{\psi\psi} = \sin^2\phi \, T_{\phi\phi}. \tag{41}$$

Like the charged blackfold, this stress tensor also have trace anomaly which shows that the corresponding lagrangian is not invariant under conformal transformation. It can be seen easily that this quasilocal stress tensor is also conserved.

4. Dual theory for neutral blackfolds

As the Brown-York method does not work for all intrinsic metric in the reference spacetime, it is not generally well defined. So in this section, we calculate the renormalized holographic stress energy tensor [27, 28] for the case of neutral blackfolds. Although holography is expected for non AdS asymptotics, such as nonrelativistic dualities, it is very well understood for asymptotically locally AdS spacetimes [1]. For every AAdS spacetime the expectation value of the dual CFT stress energy tensor is defined as:

$$< {T^{CFT}}_{ab} >= -\frac{2}{\sqrt{-g_{(0)}}}\frac{\delta S_{ren}}{\delta g_{(0)}^{ab}} \tag{42}$$

$g_{(0)}$ is the metric on the boundary and S_{ren} denotes the renormalized bulk action. In this case counterterms have been added to remove the volume divergences. We take advantages of Fefferman and Graham [29] and obtain an asymptotic solution of Einstein's field equations with conformal structure. The corresponding solution is given by,

$$ds^2 = l^2(\frac{d\rho^2}{4\rho^2}+\frac{1}{\rho}g_{ab}(x,\rho)dx^a dx^b) \tag{43}$$

where $a, b = 1, ..., D-1$, $d = D-1$. And

$$g(x,\rho) = g_{(0)} + ... + \rho^{\frac{d}{2}} g_{(d)} + h_{(d)} \rho^{\frac{d}{2}} \log\rho + ... \tag{44}$$

the coefficient $h_{(d)}$ is present only when d is even. We will follow along the same line done in [30] and determine the explicite form of the dual renormalized holographic stress tensor,

$$< {T^{CFT}}_{ab} >= \frac{dl}{16\pi G_N}({g_{(d)}}_{ab} + {X^{(d)}}_{ab}) \tag{45}$$

where ${X^{(d)}}_{ab}$ depends on the dimension. It's form for all odd d is presented in [30], and for even d it vanishes,

$${X^{(2k+1)}}_{ab} = 0 \tag{46}$$

Now we try to find the dual renormalized stress tensor for the neutral blackfolds. As we know the metric (4) is Ricci-flat and there is a very nice connection between AdS black brane (which some of it's coordinates are compactified) and Ricci-flat spacetime [31, 32]. So one can find AAdS form of that metric. the Schwarzchild black p-brane in $D = n + p + 3$ dimension has the form:

$${ds^2}_0 = -(1 - \frac{{r_0}^n}{r^n})dt^2 + \sum_{i=1}^{p} {dz_i}^2 + (1 - \frac{{r_0}^n}{r^n})^{-1} dr^2 + r^2 {d\Omega^2}_{n+1} \tag{47}$$

So the required components are given by:

$$\begin{aligned} &{ds^2}_{p+2} = -(1 - \frac{{r_0}^n}{r^n})\frac{l^2}{r^2}dt^2 + \sum_{i=1}^{p} \frac{l^2}{r^2}{dz_i}^2 + (1 - \frac{{r_0}^n}{r^n})^{-1} \frac{l^2}{r^2}dr^2 + l^2 {d\Omega^2}_{n+1}\,, \\ &\phi(\rho,x) = (n+p+1)\ln(\frac{r}{l}). \end{aligned} \tag{48}$$

By changing $n \leftrightarrow -d$, the AdS form of (47) which is a planar AdS black brane with $l = 1$ (in $d+1$ dimension) is as follows,

$${ds^2}_\Lambda = -\frac{1}{r^2}(1 - \frac{r^d}{{r_0}^d})dt^2 + \frac{1}{r^2}({dz_i}^2 + d\vec{y}^2) + \frac{1}{r^2(1 - \frac{r^d}{{r_0}^d})}dr^2, \tag{49}$$

This metric is the AdS black brane and the Fefferman-Graham coordinate (43) is the best form for this theory to find the dual stress tensor. In order to do that, we must extract the relation between r in (49) and ρ in (43). So we have,

$$\frac{1}{r^2(1-\frac{r^d}{{r_0}^d})}dr^2 = \frac{d\rho^2}{4\rho^2} \tag{50}$$

then,

$$(\frac{r}{r_0})^d = \frac{4\rho^{\frac{d}{2}}}{(\rho^{\frac{d}{2}}+1)^2} \tag{51}$$

The corresponding two important functions will be:

$$\frac{1}{r^2} = \frac{1}{r_0^2 4^{\frac{2}{d}}\rho}(\rho^{\frac{d}{2}}+1)^{\frac{4}{d}}, \tag{52}$$

$$\frac{1}{r^2}(1-\frac{r^d}{{r_0}^d}) = \frac{1}{r_0^2 4^{\frac{2}{d}}\rho}(\rho^{\frac{d}{2}}+1)^{\frac{4}{d}-2}(\rho^d - 2\rho^{\frac{d}{2}}+1), \tag{53}$$

Consequently the AdS black brane in Fefferman-Graham coordinate take the following form,

$$ds^2{}_\Lambda = -\frac{1}{r_0^2 4^{\frac{2}{d}}\rho}(\rho^{\frac{d}{2}}+1)^{\frac{4}{d}-2}(\rho^d-2\rho^{\frac{d}{2}}+1)dt^2 + \frac{1}{r_0^2 4^{\frac{2}{d}}\rho}(\rho^{\frac{d}{2}}+1)^{\frac{4}{d}}[d{z_i}^2 + d\vec{y}^2] + \frac{d\rho^2}{4\rho^2} \tag{54}$$

In the case of $D = 10 \quad \rightarrow \quad d = 9$, the asymptotically AdS metric can be brought into the following geometry near the boundary ($\rho = 0$),

$$ds^2{}_\Lambda = -\frac{1}{r_0^2 4^{\frac{2}{9}}\rho}(1-\frac{32}{9}\rho^{\frac{9}{2}}+\frac{494}{81}\rho^9+...)dt^2 + \frac{1}{r_0^2 4^{\frac{2}{9}}\rho}(1+\frac{4}{9}\rho^{\frac{9}{2}}-\frac{10}{81}\rho^9+...)[d{z_i}^2+d\vec{y}^2] + \frac{d\rho^2}{4\rho^2} \tag{55}$$

By using the relation (44) one can obtain the explicit form of the metric coefficients which are given by,

$$g_{(0)} = \frac{1}{r_0^2 4^{\frac{2}{9}}}diag(-1,1,...,1), \tag{56}$$

$$g_{(9)} = \frac{1}{r_0^2 4^{\frac{2}{9}}}diag(\frac{32}{9},\frac{4}{9},...,\frac{4}{9}), \tag{57}$$

Finally the dual stress energy tensor in terms of relation (45) is

$$< T^{CFT}{}_{ab} > = \frac{9}{16\pi G_N}({g_{(9)}}_{ab} + X^{(9)}{}_{ab}) = \frac{9}{16\pi G_N}{g_{(9)}}_{ab} \ . \tag{58}$$

This stress tensor can be checked formally as a conserved and traceless quantity. It is traceless because the conformal anomaly evaluated for global AdS vanishes. If the metric (55) be locally boosted, it takes the form,

$$ds^2 = -\frac{1}{r_0^2 4^{\frac{2}{9}} \rho}(4\rho^{\frac{9}{2}} - \frac{504}{81}\rho^9 + ...)u_a u_b dx^a dx^b + \frac{1}{r_0^2 4^{\frac{2}{9}} \rho}(1 + \frac{4}{9}\rho^{\frac{9}{2}} - \frac{10}{81}\rho^9 + ...)[\eta_{ab} dx^a dx^b + d\vec{y}^2] + \frac{d\rho^2}{4\rho^2} \tag{59}$$

Then,

$$g_{(9)ab} = \frac{1}{r_0^2 4^{\frac{2}{9}}}(4u_a u_b, \frac{4}{9}\eta_{ab}, ..., \frac{4}{9}\eta_{ab}), \tag{60}$$

So the renormalized holographic stress tensor takes the general representation:

$$< T^{CFT}{}_{ab} >= \frac{9}{16\pi G_N}[\frac{1}{r_0^2 4^{\frac{2}{9}}}(4u_a u_b + \frac{4}{9}\eta_{ab})] \ . \tag{61}$$

The boundary metric is conformally flat and the extracted stress energy tensor takes the ideal conformal fluid form in 9-dimensional spacetime. Comparison of (61) by (9) concludes the pressure and energy density of this blackfold such that:

$$p = \frac{1}{4^{\frac{11}{9}} \pi G_N r_0^2}, \qquad \varepsilon = \frac{8}{4^{\frac{11}{9}} \pi G_N r_0^2} \ . \tag{62}$$

It also proves that the resulted stress tensor (which determine the boundary fluid and comes from the duality of gravity and fluid) is conserved and traceless.

For non AAdS spacetime we can not use the represented method, because the holographic principle is not very well understood for them. Hopeful this method be extended such that it be consistent for more spacetimes.

5. Conclusion and outlook

In the covariant theory, the local energy momentum density of the gravitational field is not prevalent. Instead, a quasilocal stress tensor can be defined on the boundary of spacetime. But resulting stress tensor may diverge when the boundary goes to infinity. In order to solve this difficulty one can add an appropriate boundary term to the action which is completely relevant.

Brown and York proposes a method to remove divergences. But it does not work for all intrinsic metric in the reference spacetime, so the Brown-York procedure is not generally well defined. Fortunately in AAdS spacetimes one can overcome to this problem. As gravitational action of the bulk corresponds to the quantum effective action of a conformal field theory on the AdS boundary, the expectation value of the stress tensor in the CFT solves the problem.

In this paper we considered the metric of charged and neutral blackfold and obtained the boundary stress tensor. As the induced Brown-York stress tensor is conformally flat, the holographic stress tensor should be conformal to Brown-York stress energy tensor. For the case of neutral blackfold which it's metric is Ricci-flat, by applying the AdS/Ricci-flat correspondence, we fined the corresponding AAdS form for it. Then for $D = 10$ the renormalized holographic boundary stress tensor extracted. As we expected, the results show that the corresponding stress tensor are conserved and traceless. In that case there is not any conformal anomaly for the pointed theory that proves the lagrangian of the neutral blackfold is invariant under conformal transformation. Also it takes the form of ideal fluid with determined energy density and pressure.

Acknowledgment

The authors would like to thank Borzoo Nazari for helpful discussion and comments in the subject of Brown-York stress energy tensor.

References

[1] Kostas skenderis, Lecture Notes on Holographic Renormalization, Class.Quant.Grav. 19:5849-5876 (2002), [arXiv:0209067].

[2] Juan M. Maldacena, The Large N Limit of Superconformal Field Theories and Supergravity , Adv.Theor.Math.Phys. 2:231-252 (1998), [arXiv:9711200].

[3] S.S. Gubser, I. R. Klebanov, A. M. Polyakov, Gauge Theory Correlators from Non-Critical String Theory , Phys.Lett.B428:105-114, (1998), [arXiv:9802109].

[4] O. Aharony, S. S. Gubser, J. Maldacena, H. Ooguri, Y. Oz, Large N Field Theories, String Theory and Gravity , Phys.Rept.323:183-386, (2000), [arXiv:9905111].

[5] Edward Witten, Anti De Sitter Space And Holography , Adv.Theor.Math.Phys.2:253-291, (1998), [arXiv:9802150].

[6] Sayantani Bhattacharyya, Veronika E Hubeny, Shiraz Minwalla, Mukund Rangamani, Nonlinear Fluid Dynamics from Gravity , JHEP 0802:045, (2008), [arXiv:0712.2456].

[7] Veronika E Hubeny, The Fluid/Gravity Correspondence: a new perspective on the Membrane Paradigm , Class.Quant.Grav.28:114007, (2011), [arXiv:1011.4948].

[8] Natalia Pinzani-Fokeeva, Marika Taylor Towards a general fluid/gravity correspondence , Phys. Rev. D 91, 044001 (2015), [arXiv:1401.5975].

[9] Robrto Emparan, Troels Harmark, Vasilis Niarchos, Niels A. obers, World-Volume Effective Theory for Higher-Dimensional Black Holes , Phys.Rev.Lett.102:191301, (2009).[arXiv:0902.0427].

[10] Robrto Emparan, Troels Harmark, Vasilis Niarchos, Niels A. obers,Maria J. Rodrigues, The Phase Structure of Higher-Dimensional Black Rings and Black Holes , JHEP 0710:110, (2007).[arXiv:0708.2181].

[11] Robrto Emparan, Troels Harmark, Vasilis Niarchos, Niels A. obers, Essentials of Blackfold Dynamics , JHEP 1003:063, (2010).[arXiv:0910.1601].

[12] Justin R. David, Manavendra Mahato, Spenta R. Wadia Hydrodynamics from the D1-brane , JHEP 0904:042, (2009).[arXiv:0901.2013].

[13] Ingnar Kanitscheider, Kostas Skenderis, Universal hydrodynamics of non-conformal branes , JHEP 0904:062, (2009).[arXiv:0901.1487].

[14] Geoffrey Compere, Paul McFadden, Kostas Skenderis, Marika taylor, The relativistic fluid dual to vacuum Einstein gravity , JHEP 03(2012)076 (2012).[arXiv:1201.2678].

[15] Christopher Eling, Adiel Mayer, Yaron Oz, The Relativistic Rindler Hydrodynamics , [arXiv:1201.2705].

[16] Irene Bredberg, Andrew Strominger, Black Holes as Incompressible Fluids on the Sphere , ,[arXiv:1106.3084].

[17] Tai-Zhuo Huang, Yi Ling, Wen-Jian Pan, Yu Tian, Xiao-Ning Wu, From Petrov-Einstein to Navier-Stokes in Spatially Curved Spacetime , JHEP 1110:079, (2011).[arXiv:1107.1464].

[18] Ryuichi Nakayama, The Holographic Fluid on the Sphere Dual to the Schwarzschild Black Hole , Publisher (year).[arXiv:1109.1185].

[19] Dionysios Anninos, Tarek Anous, Irene Bredberg, Gim Seng Ng, Incompressible Fluids of the de Sitter Horizon and Beyond , [arXiv:1110.3792].

[20] Chao Niu, Yu Tian, Xiao-Ning Wu, Yi Ling, Incompressible Navier-Stokes Equation from Einstein-Maxwell and Gauss-Bonnet-Maxwell Theories , Phys.Lett.B711:411-416 (2012), [arXiv:1107.1430].

[21] Cheng-Yong Zhang, Yi Ling, Chao Niu, Yu Tian, Xiao-Ning Wu, Magnetohydrodynamics from gravity , Phys. Rev. D 86, 084043 (2012), [arXiv:1204.0959].

[22] Vyacheslav Lysov, On the Magnetohydrodynamics/Gravity Correspondence , [arXiv:1310.4181].

[23] Brandon Carter, Essentials of Classical Brane Dynamics , Int.J.Theor.Phys. 40 2099-2130 (2001), [arXiv:0012036].

[24] J. David Brown, James W. York, Quasilocal Energy and Conserved Charges Derived from the Gravitational Action , Phys.Rev.D 47:1407-1419, (1993), [arXiv:9209012].

[25] Roberto Emparan, Troels Harmark, Vasilis Niarchos, Niels A. Obers, Blackfolds in Supergravity and String Theory , JHEP 1108 154 (2011), [arXiv:1106.4428].

[26] Vijay Balasubramanian, Per Kraus, A Stress Tensor for Anti-de Sitter Gravity , Commun.Math.Phys. 208 413-428 (1999), [arXiv:9902121].

[27] Mans Henningson, Kostas Skenderis, The Holographic Weyl anomaly , JHEP 9807 023 (1998), [arXiv:9806087].

[28] Sebastian de Haro, Kostas Skenderis, Sergey N. Solodukhin, Holographic Reconstruction of Spacetime and Renormalization in the AdS/CFT Correspondence , Commun.Math.Phys. 217 595-622 (2001), [arXiv:0002230].

[29] C. Fefferman, C. Robin Graham, Elie Cartan et les Mathe'matiques d'aujourd' hui , (Aste'risque, 1985) 95,

[30] Kostas Skenderis, Asymptotically Anti-de Sitter spacetimes and their stress energy tensor , Int.J.Mod.Phys. A 16 740-749 (2001), [arXiv:0010138].

[31] Marco M. Caldarelli, Joan Camps, Blaise Goutraux, Kostas Skenderis, AdS/Ricci-flat correspondence and the Gregory-Laflamme instability , Phys. Rev. D 87, 061502(R) (2013), [arXiv:1211.2815].

[32] Marco M. Caldarelli, Joan Camps, Blaise Goutraux, Kostas Skenderis, AdS/Ricci-flat correspondence , JHEP 04 071 (2014), [arXiv:1312.7874].